

Original Research Article

A DEXA scan at 20 years of age in select subjects may show a z-score "shift to the left" giving time after ethnic corrections to regain optimal peak BMD at 30 years

Kiran Kumar Venkata Satya Nandivada*, Adithya Sridhar Acharya, Kabilesh Jothilingam, Shabbeer Imtiaz Ahamed, Ajin Mathews John, Vaishnavi Nandivada, Aysha Naushad Khan, Adnan Mirza, Usama Ahamed Rizwi, Maliha Karim, Chandana Kalapala, Sagarika Jayakumar, Praharsha Pothina, Ishan Mohzin

International Research Trainee Academy, Seethammadhaa, Visakhapatnam, Andhra Pradesh, India

Received: 27 September 2026

Revised: 12 October 2026

Accepted: 15 October 2026

*Correspondence:

Dr. Kiran Kumar Venkata Satya Nandivada,

E-mail: kows.foru@gmail.com

Copyright: © the author(s), publisher and licensee Medip Academy. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Background: Osteoporosis is a silent pandemic, and it's vital to detect osteopenia early. This study shows the importance of a DEXA baseline scan at 20 years of age as a part of a PREVENTIVE MEASURE to gain time to reach the optimum Peak BMD at around 30 years.

Methods: Among 5,000 patients tested in a mixed or ambispective study determining BMD between 2000-2026, by the IRT programme by NGO KRUSHI OWS.

The exclusion criteria were patients with known causes of bone weakness due to congenital diabetes, Thyroid deficiencies, renal dysfunction, familial history of bone weakness.

Results: Osteopenia was prevalent in 60% (n=3,000), Osteoporosis was found in 15% (n=750), and Normal Bone Density was found in 25% (n=1,250). The analysis showed that 75% of study patients had low bone density, highlighting the consequences of low bone mass for individuals, families, and insurance companies.

Conclusions: If the osteopenia stage, a precursor to osteoporosis, is missed or neglected, the opportunity to reach optimal peak BMD is lost forever, making this study an important call for attention from osteoporosis foundations worldwide.

Keywords: Osteopenia, Osteoporosis, DEXA scan, T score, Z score

INTRODUCTION

Osteopenia is a precursor of osteoporosis, and if diagnosed with precision, it can be found in growing fetuses, children, and young adults, but sadly it is often unrecognised and uninvestigated. Peak bone mass is generally achieved by the 3rd decade of life and inadequate attainment of PBMD predisposes to osteopenia and later osteoporosis; if it occurs in the young, the susceptibility to fragility fractures increases even in adolescence or early

adulthood, as the rapid growth of the bone with lagging bone density leads to a 'transient deficiency of cortical bone mass in growing adolescents or a latent physiological osteopenia of growing bones during adolescents'.¹⁻⁴ Also, in this case report by Sheikdon et al transient osteoporosis of the hip was reported in a 17-year-old female.⁵ The objective of this study is to identify low Z-scores early in young adulthood and rectify them before peak BMD is achieved.

A low Z score or low bone mineral density should alert the clinician to start timely correction so the patient reaches peak BMD by 25 to 30 years, after which bone density falls gradually in both genders, though with different plateau and decline patterns.

METHODS

This is a mixed or ambispective study done by the Research Centre of KRUSHI Orthopaedic Welfare Society, an NGO based in Andhra Pradesh, India, between April 2000 and 2026. Statistical analysis, including data collection, organisation, analysis, interpretation, and presentation, was done. The Selection criteria included 5000 screened patients at the KOWS research centre and 200 articles from databases such as PubMed.

This study emphasises the importance of using quantitative ultrasound BMD-measuring equipment in mass screening camps as a screening tool only and, in selected patients, DEXA (Dual Energy X-ray Absorptiometry) for confirmation, if available. All participants provided informed consent. Patient privacy was protected, and the study adhered to the Helsinki guidelines. Ethical approval was obtained from the Krushi Orthopaedic Welfare Society Ethics Committee, Visakhapatnam, Andhra Pradesh, India.

RESULTS

The Demographic data focused on both males and females around 20 years of age, Indian Ethnicity, students studying in tribal, rural and urban colleges without income, unmarried and non-religion-centric, with nuclear families, with normal-appearing health status with symptoms of fatigue, bone and joint pain, fractures following minor trauma, vertebral pain and normal inflammatory markers.




Figure 1: Pie Diagram showing IRT programme data from the osteopenia study conducted between 2000 and 2026, showing normal BMD in only 25% of young adults before 30 years of age, osteopenia at 60%, and Osteoporosis at 15% with a history of frequent fractures.

Osteopenia was prevalent in 60% (n=3,000), Osteoporosis was found in 15% (n=750), and normal bone density was found in 25% (n=1,250).

The analysis showed that 75% of study patients had low bone density, highlighting the consequences of low bone mass for individuals, families, and insurance companies.




Figure 2: The demographic, gender distribution of the findings in this study are displayed above.

Osteopenia was prevalent in 60% (n=3,000), Osteoporosis was found in 15% (n=750), and Normal Bone Density was found in 25% (n=1,250).

DISCUSSION

Preventive orthopaedics is important in osteoporosis, as it can completely disable a person at any age. The analysis showed that 75% of study patients had low bone density, highlighting the consequences of low bone mass for individuals, families, and insurance companies.

In the article by Panchal et al in the *Journal of Preventive Medicine and Holistic Health* 2025;11(1):28-37, osteopenia in young adults across India was emphasised as Osteopenia (43.91%) and Osteoporosis (8.40%), out of a total of 5,000 subjects. Our 25-year study has similar findings and is the reason for this research publication. Poor lifestyles, smoking during pregnancy or in regular life, alcoholism, poor diet with low vitamin D levels and lack of physical activity during pregnancy appear to be detrimental to bone mineral deposition during intrauterine life.

Growth hormone deficiency is reported to cause Higher fracture risk in COGHD (childhood-onset GH deficiency), and treatment with GH should minimise osteoporosis in adulthood. Further, optimum PEAK BMD is reached by an optimum GH level at the age of transition to young adulthood by treatment with rhGH in growing children with COGHD.⁶

Trabecular bone loss is seen in patients with Collagen VI α 2 chain deficiency by increasing osteoclast differentiation by activating the TNF α signalling pathway.⁷ Collagen supplements are more beneficial when taken as an adjuvant with Vitamin D and Calcium supplements.⁸ Osteogenesis imperfecta is another common genetic condition related to type 1 collagen.

In utero fetal osteopenia or Osteoporosis does exist and can result from a metabolic bone disease linked with premature birth due to reduced transfer of calcium and phosphorus in the third trimester, or a particular genetic aberration

causing demineralisation in the fetus. Once delivered, the newborn may show low birth weight, which is usually caused by maternal low Vitamin D; maternal illness, lack of breast milk or refusal to breastfeed can lead to osteopenia in the newborn or the infant if proper substitute feed is not given.⁹⁻¹²

Ethnic BMD variations exist, as seen in Caucasian bones, which are stronger than those of Asians. Hence, the Hologic data, when applied to Asians, gives more false positives, as Asians are reported as having a higher incidence of osteopenia or osteoporosis, and this led the ICMR to release conversion formulas to rectify this error in the T and Z scores. This IRTIP research is supported by a publication by Babhulkar et al which concluded that osteopenia in younger populations was more prevalent and that enough awareness to detect it was absent and called for pan-India studies in this regard, which is also a supportive lead to the present study.¹³

The highest BMD was found in the Tobagonian African race and is due to genetic factors (Genes VDR, ER α , and COL1A1), which show 10–20% higher bone mineral density at the hip and spine, with thicker cortical bone and stronger trabecular structure contributing to higher bone mass and lower resorption rates.¹⁴ The hormonal influence resulting in a “Vitamin D paradox” is an interesting condition characterised by efficient calcium conservation and enhanced skeletal sensitivity to parathyroid hormone.¹⁵

The observation of ‘transient deficiency of cortical bone mass in growing adolescents’ is important because, during active growth, peak bone mineral density reached only 57% of the total bone mineral content while the height velocity grows to 90% of expected adult height and there is more transient weakness in the forearms which may fracture on trivial injuries and this study highlights this Preventive orthopaedics aspect.¹⁶

After linear growth is reached the bone continues to remodel at a rate of up to 50% per year. Bone remodelling is a delicate balance between resorption and formation and is regulated by PTH, vitamin D, insulin-like growth factor, and calcitonin. The male skeleton has greater resistance to mechanical stress due to higher BMD than in girls, and in boys the appendicular skeleton shows predominant periosteal deposition, making the cortical layers thicker as the periosteal and endosteal surfaces are thickened.

The same increased cortical thickness is found in the vertebrae of the axial skeleton. In girls, endosteal deposition is seen predominantly. Sexual structural dimorphism presents as a 10-15% increase in the frontal bone in men compared with women, providing greater mechanical resistance to trivial fractures in males.¹⁷ It is prudent to understand that, from bone formation onward, females are susceptible to trivial injuries and may sustain fractures and failure patterns. The DEXA machine uses dual-energy X-ray absorptiometry and is very accurate,

though not mobile at this time. The appearance of a BMD Curve differs in males and females.¹⁸

Safety of DEXA scan: DEXA scan, or dual-energy X-ray absorptiometry, is very safe, as the radiation dose given by a DEXA scan to a patient is usually only 0.001–0.01 mSv (1–10 μ Sv) (equal to 100 chest X-rays). It's better to test for the bone density with a DEXA SCAN rather than overlook a silent disease with progressive weakening of the axial and appendicular skeleton. Comparison with similar findings in support of this study.




Figure 3 : Different Peak bone mass densities in different individuals are depicted in different genders, where the PBMD reached is higher in males with a higher plateau and then gradually falls, while in females there is a lower PBMD with a flatter, shorter plateau, which suddenly declines after menopause, and at the bottom there is an interrupted curve in either gender where the PBMD is seldom reached to the expected levels and remains in deficit throughout and also declines into insufficiency, earlier osteopenia and earlier severe osteoporosis.

This is why females are always in deficit in their BMD, as the peak BMD is never as high as in males, and they will lose more bone density much earlier; if a 10% higher Peak BMD is achieved, the onset of osteoporosis is delayed by 10 years.¹⁹ The problem in India is that the bones are less dense than Caucasian bones of the Europeans, which were analysed by the NHANES-III database, and when the caucasian data was used to calibrate the Hologic machines for inventing the T score and the Z SCORE and then used for the Indian population, the T and Z scores were lower, which were normal for Indians.²⁰ This prompted the Indian Council of Medical Research to conduct a retrospective study called ‘Ethnica’, where special conversion formulas were developed to get the reliable ICMR (T score) and the ICMR (Z score) and the false positives or over declaration of osteopenia and osteoporosis was eliminated when the ICMR data were incorporated into the software systems of the hologic DEXA and ultrasound machines to avoid drug-induced organ overload and failure.²¹ For this in India, the Indian Council of Medical Research (ICMR) provided

rough ranges as reference ranges; the T-scores increase by roughly +0.51 (HIP) and by +1.64 (Spine).

Shetty et al used the Indian database from 4 Indian cities versus the Hologic database in postmenopausal women and they found that a high incidence of false-positive osteoporosis when the Hologic Caucasian data was used was in fact osteopenia when the Indian databases were incorporated into the software.²² This information necessitates the development of Indian BMD machines for our 1.5 billion Indians as part of the 'Make in India' initiative to assist in early detection, as India is one of the leading countries with the highest number of osteopenic and osteoporosis patients in the world. In the article by Kanis et al which highlights the term "Gradient of risk for hip fracture," the authors discuss the prognosis for individuals. If it is 2.6, this indicates that the risk of hip fracture increases by 2.6 times for every 1 SD reduction in hip BMD.²³ The Chinese also showed the same discrepancy like in Indians, which was portrayed in a study by Wang J et al wherein the Chinese women showed a discrepancy in the T score when compared to the Italian women (-3.75 = -2.44), pointing to an overdiagnosis of osteoporosis in the Chinese.²⁴ Hence, in India or China, this kind of discrepancy gets amplified as these countries have the world's largest populations. Nutritional deficiencies in calcium and Vitamin D in developing countries like India also adversely contribute to bone health. Bone mineral density in the young, as per the WHO, is reported as a Z score, which is the BMD in comparison to the mean BMD for peers at the same age, gender and race and as a T score, which is the BMD in comparison to the expected T score compared to the peak BMD value at around 30 years of age.

One would expect to see the best bone mineral density for oneself, and this, at around forty years of age, may look like the report below




Figure 4: Shows the best BMD T-score of +3.7 in a 45-year-old male patient.

The picture emphasises the best BMD for age, but at the same time, in another individual below, the T-score may

show a notable "Shift to the left", and it means that osteopenia or osteoporosis already exists, and this is a red flag for urgent intervention, as seen in our free BMD camps since 2000.

Two scientific papers were presented in 2003 at the APIOACON and IOACON conferences in India to highlight the point that osteopenia must be detected as early as possible to reduce the risk of untreated osteoporosis developing, at a time when little research was highlighting this aspect and is being continued as COE on Online lectures in clernet.com and IJCR Central platforms. Enclosed below are two certificates as evidence of this, stressing the need for mass screening for osteopenia and osteoporosis since 2001, i.e., for twenty-five years.

Formulas that one must know when dealing with Caucasian data and ICMR databases: Body mass index: Weight (kg) / height in meter² was used to calculate body mass index (BMI). T Score (Standard): BMD of subject (gms/cms²) - Ref BMD / SD (G/cm²). Formula for T score [ICMR]: BMD subject - BMD ref / SD ref (ICMR). Formula for Z score [ICMR]: BMD(Subject)- BMD ref / Standard deviation ref ICMR.

IRT programme wishes to highlight a DEXA scan with T-score shifted to the left, a RED FLAG in DEXA SCANS. Presentation of actual data collected by the IRT programme from some patients:




Figure 5: A DEXA scan report of a male patient of 20 years of age from Visakhapatnam, Andhra Pradesh, India showing a T score of -5.6, meaning severe OSTEOPOROSIS, a classic example of a T score that has shifted to the left, which implies that on a trivial injury, his fracture risk is expected to be 15 to 20 times higher than that of a normal man of his age. [Even if it is corrected by the rough guide of + 1.5 to this T score to suit our Indian patient, the corrected score comes to -4.1, and if the Z score has to be evaluated, then it can be evaluated using the ICMR FORMULA].

We have 2 options for checking one's T-score and Z-score.

The first is an easier piece of equipment to handle and move around: a non-radiation-based modality called quantitative Ultrasound bone mineral density screening. In 2023, an article by Kushwaha NS et al. also validates the usage of QUS equipment for screening osteopenia and osteoporosis.²⁶ This study from 2000 is supported by the Osteolaus study by Métrailler A et al which supports the use of the heel Achilles ultrasound machine as a QUS combined with clinical risk factors (CRFs) as a pre-screening tool in spaces where the DEXA scan is not available, but it was concluded that it cannot be used for further monitoring.²⁷

A DEXA scan is institutional, immovable, and highly expensive, but sensitive and specific to 99%, with a radiation exposure far less than a chest X-ray. Now, the latest tool is the Trabecular Bone Score. FRAX score with ethnic adjustments, along with other assessment tools, are indeed helpful.

We should consider BMD measurement in young patients complaining of fatigue, bone and joint pains, thoracolumbar aches, falls and muscle weakness, loss of height or spinal deformity early in life, and an increase in the scoliosis curves because if we do not recognise the weakness in the bone density, precious time to rectify is lost permanently before the peak bone mass density is reached.

In an article by C. J. Hernandez et al it is noted that a 10% increase in peak bone mass density can delay the onset of osteoporosis by 13 years.²⁸

In the foetus, infantile osteoporosis is a real entity. In the article by Marvin Miller, the UTAH paradigm emphasises that foetal movements reflect foetal bone strength and that poor mineralisation in infants may also seem to be due to child abuse.²⁹

Multifactorial bone fragility disorders in young infants could be due to the following; Genetic causes: Mucopolysaccharidosis (MPS)³⁰, Osteogenesis Imperfecta³¹; Metabolic Causes: Low Vitamin D³²; Congenital rickets - X-linked hypophosphatemia³³; Maternal habits: non-active lifestyle, low maternal vitamin levels, along with other deficiencies, Thyroid abnormalities, diabetes and medications.

Dual-energy X-ray absorptiometry (DXA) is used to evaluate bone health, though interpreting results in infants' children with severe short stature and deformities is challenging. In the 2023 article by Ciancia et al Osteoporosis in children and adolescents was described, and anti-resorptive medications were identified as the first-line treatment.³⁴

In Idiopathic Scoliosis the Curve severity has an inverse relationship with bone mineral mass during peri-puberty in adolescent Idiopathic Scoliosis, suggesting that prevention

of osteopenia could play a role in controlling spinal progression.^{35,36}

Interpretation of BMD DEXA scans in children is challenging as the low or high Z scores stare at the viewer [Osteopetrosis][Fluorosis] and the meaning of a Z score must be taken into consideration.

Bone Mineral Density (BMD) discordance: a DEXA scan often shows osteopenia or even osteoporosis appearing in only one location, i.e., in the Lumbar Spine. At the same time, the distal radius or the neck of the femur has normal or osteopenic T-scores [Bone Mineral Density (BMD) discordance].³⁷ This is BMD Discordance that confuses the clinician to choose just exercises, nutrients, vitamin D and Elemental calcium, or bisphosphonates, to be introduced and followed up. The incidence of fragility fractures is higher in the Osteopenic group than in the osteoporotic group.³⁸

Low testosterone levels due to hypogonadism can cause osteopenia and need therapeutic correction to restore BMD in time and can cause atherosclerotic plaques, calcification, and osteoporosis later as the individual ages.^{39,40} Yet TRT, or testosterone replacement therapy in males with osteopenia, is still inconclusive as of today, with larger trials underway. In osteoporosis, the upper end plates are fractured more than the lower end plates (Men > women).⁴¹

In addition to a better lifestyle with weight-bearing exercises, avoidance of smoking, alcohol, and medications in selective cases, an emerging Therapy is a non-pharmacological option, OSTEOBOOST – a kind of vibration therapy.⁴²

A newer tool called REMS, or Radiofrequency Echographic Multi Spectrometry (REMS), is being tested for diagnosing osteoporosis in a European multicentre trial.⁴³ As per Mobasseri M et al for monitoring osteopenia, evaluating bone turnover markers have a significant relationship with lower BMD.⁴⁴

CONCLUSION

In this 25 years research study of young 20 years aged patients suggestive of osteopenia with normal inflammatory markers, vertebral pain, bone and joint pains - early intervention with a DEXA scan for screening for osteopenia is the concluding suggestion of this study as this disease is a silent pandemic without clear cut clinical or laboratory findings and only a high clinical suspicion assists the young patients to reach their Peak Bone Mineral Density before 30 years after which the density is by nature designed to decline and cause further trivial fractures, loss of vertebral height, receding gums, loss of teeth and stamina -The osteoporosis sequelae. This study highlights that 60% of the screened youngsters showed osteopenia and 15 % showed osteoporosis, meaning the detection was actually already late for the best results from nonmedical

and medical treatment and that even at this stage early intervention could help them achieve their best Peak BMD.

Funding: No funding sources

Conflict of interest: None declared

Ethical approval: The study was approved by the Institutional Ethics Committee

REFERENCES

1. Xue S, Kemal O, Lu M, Lix LM, Leslie WD, Yang S. Age at attainment of peak bone mineral density and its associated factors: the National Health and Nutrition Examination Survey 2005–2014. *Bone*. 2020;131:115163.
2. Chevalley T, Rizzoli R. Acquisition of peak bone mass. *Best Pract Res Clin Endocrinol Metab*. 2022;36(2):101616.
3. Weaver CM, Gordon CM, Janz KF, Kalkwarf HJ, Lappe JM, Lewis R, et al. The National Osteoporosis Foundation's position statement on peak bone mass development and lifestyle factors: a systematic review and implementation recommendations. *Osteoporos Int*. 2016;27(4):1281–386.
4. Wang Q, Wang X, Iuliano-Burns S, Ghasem-Zadeh A, Zebaze R, Seeman E. Rapid growth produces transient cortical weakness: a risk factor for metaphyseal fractures during puberty. *J Bone Miner Res*. 2010;25(7):1521–6.
5. Sheikdon AA, Ahmed AM, Ahmed OH. Transient osteoporosis of the hip in a 17-year-old nonpregnant female: a case report. *J Med Case Rep*. 2025;19(1):323.
6. Doknic M, Stojanovic M, Markovic A. Transition period and young adulthood in patients with childhood onset growth hormone deficiency (COGHD): impact of growth hormone replacement on bone mass and body composition. *Int J Mol Sci*. 2024;25(19):10313.
7. Pham HT, Kram V, Dar QA, Komori T, Ji Y, Mohassel P, et al. Collagen VI α 2 chain deficiency causes trabecular bone loss by potentially promoting osteoclast differentiation through enhanced TNF α signaling. *Sci Rep*. 2020;10(1):13749.
8. Sun C, Yang A, Teng F, Xia Y. Efficacy of collagen peptide supplementation on bone and muscle health: a meta-analysis. *Front Nutr*. 2025;12:1646090.
9. Miller M, Hangartner T. Temporary brittle bone disease: association with decreased fetal movement and osteopenia. *Calcif Tissue Int*. 1999;64:137–143.
10. So KW, Ng PC. Treatment and prevention of neonatal osteopenia. *Curr Paediatr*. 2005;15(2):106–13.
11. Perrone S, Caporilli C, Grassi F, Ferrocino M, Biagi E, Dell'Orto V, et al. Prenatal and neonatal bone health: updated review on early identification of newborns at high risk for osteopenia. *Nutrients*. 2023;15(16):3515.
12. Tabatabaei RS, Mirjalili SR, Ashrafi A, Shamsi F. The relationship between maternal vitamin D levels and osteopenia development in preterm infants: a cross-sectional study. *Int J Reprod Biomed*. 2024;22(11):845–56.
13. Babhulkar S, Seth SS. Prevalence of osteoporosis in India: an observation of 31238 adults. *Int J Res Orthop*. 2021;7:362–8.
14. Zmuda JM, Yerges LM, Kammerer CM, Cauley JA, Wang X, Nestlerode CS, et al. Association analysis of WNT10B with bone mass and structure among individuals of African ancestry. *J Bone Miner Res*. 2009;24(3):437–47.
15. Sosa-Henríquez M, Torregrosa-Suau Ó, Gómez de Tejada-Romero MJ, Cancelo-Hidalgo MJ, Tarazona-Santabalbina FJ, Etxebarria-Foronda I, et al. Rethinking vitamin D deficiency: controversies and practical guidance for clinical management. *Nutrients*. 2025;17(22):3573.
16. Akhiiarova K, Khusainova R, Minniakhmetov I, Mokrysheva N, Tyurin A. Peak bone mass formation: modern view of the problem. *Biomedicines*. 2023;11(11):2982.
17. Gilsanz V, Wren TAL, Ponrartana S, Mora S, Rosen CJ. Sexual dimorphism and the origins of human spinal health. *Endocr Rev*. 2018;39:221–39.
18. Hereford T, Kellish A, Samora JB, Reid Nichols L. Understanding the importance of peak bone mass. *J Pediatr Orthop Soc North Am*. 2024;7:100031.
19. Cherian KE, Kapoor N, Asha HS, Thomas N, Paul TV. Influence of different reference databases on categorization of bone mineral density: a study on rural postmenopausal women from southern India. *Indian J Endocrinol Metab*. 2018;22(5):579–83.
20. Bhudhikanok GS, Wang MC, Eckert K, Matkin C, Marcus R, Bachrach LK. Differences in bone mineral in young Asian and Caucasian Americans may reflect differences in bone size. *J Bone Miner Res*. 1996;11(10):1545–56.
21. Kandasamy N, Karjala SL, Jayakumar D, Machiraju PK, Kalathur HK, Balakrishnan L. Estimation of T scores with Hologic using Native vs. Caucasian data in Indians (ETHNICA): a single center retrospective study. *BMC Musculoskelet Disord*. 2025;26(1):448.
22. Shetty S, Kapoor N, Naik D, Asha HS, Thomas N, Paul TV, et al. The impact of the Hologic vs. the ICMR database in diagnosis of osteoporosis among South Indian subjects. *Clin Endocrinol (Oxf)*. 2014;81:519–22.
23. Kanis JA, McCloskey EV, Johansson H, Oden A, Melton LJ III, Khaltayev N. A reference standard for the description of osteoporosis. *Bone*. 2008;42(3):467–75.
24. World Health Organization. Assessment of fracture risk and its application to screening for postmenopausal osteoporosis. Geneva: World Health Organization; 1994. Available from: https://fraxplus.org/sites/frax/files/pdf/WHO_Technical_Report.pdf. Accessed on 11 September 2026.

25. Hadadi I. Comparative analysis of bone density measurement techniques: a systematic review of quantitative ultrasound and dual-energy X-ray absorptiometry. *Front Endocrinol (Lausanne)*. 2026;17:1768327.
26. Kushwaha NS, Singh A, Kumar S, Kumar D, Bharat A. Validation of quantitative ultrasonography for osteoporosis diagnosis in postmenopausal women compared to dual-energy X-ray absorptiometry (DEXA). *Cureus*. 2023;15(5).
27. Métrailler A, Hans D, Lamy O, Gonzalez Rodriguez E, Shevroja E. Heel quantitative ultrasound (QUS) predicts incident fractures independently of trabecular bone score (TBS), bone mineral density (BMD), and FRAX: the OsteoLaus Study. *Osteoporos Int*. 2023;34(8):1401–9.
28. Hernandez CJ, Beaupre GS, Carter DR. A theoretical analysis of the relative influences of peak BMD, age-related bone loss and menopause on the development of osteoporosis. *Osteoporos Int*. 2003;14:843–7.
29. Miller M. Foetal bone loading is a critical determinant of foetal and young infant bone strength. *Clin Biomech (Bristol, Avon)*. 2025;125:106513.
30. Jiang Z, Byers S, Casal ML, Smith LJ. Failures of endochondral ossification in the mucopolysaccharidoses. *Curr Osteoporos Rep*. 2020;18(6):759–73.
31. Subramanian S, Anastasopoulou C, Viswanathan VK. Osteogenesis imperfecta. In: StatPearls. Treasure Island (FL): StatPearls Publishing; 2026. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK536957/>. Accessed on 11 August 2026.
32. Mahon P, Harvey N, Crozier S, Inskip H, Robinson S, Arden N, et al.; SWS Study Group. Low maternal vitamin D status and fetal bone development: cohort study. *J Bone Miner Res*. 2010;25(1):14–9.
33. Ackah SA, Imel EA. Approach to hypophosphatemic rickets. *J Clin Endocrinol Metab*. 2023;108(1):209–20.
34. Ciancia S, Högl W, Sakkars RJB, et al. Osteoporosis in children and adolescents: how to treat and monitor? *Eur J Pediatr*. 2023;182:501–11.
35. Lee WT, Cheung CS, Tse YK, Guo X, Qin L, Lam TP, et al. Association of osteopenia with curve severity in adolescent idiopathic scoliosis: a study of 919 girls. *Osteoporos Int*. 2005;16(12):1924–32.
36. Bartynski WS, Heller MT, Grahovac SZ, Rothfus WE, Kurs-Lasky M. Severe thoracic kyphosis in the older patient in the absence of vertebral fracture: association of extreme curve with age. *AJNR Am J Neuroradiol*. 2005;26(8):2077–85.
37. Yoon BH, Kim DY. Discordance between hip and spine bone mineral density: a point of care. *J Bone Metab*. 2021;28(4):249–51.
38. Varacallo MA, Seaman TJ, Jandu JS, et al. Osteopenia. In: StatPearls. Treasure Island (FL): StatPearls Publishing; 2026. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK499878/>. Accessed on 11 August 2026.
39. Deb P, Gupta SK, Godbole MM. Effects of short-term testosterone replacement on areal bone mineral density and bone turnover in young hypogonadal males. *Indian J Endocrinol Metab*. 2012;16(6):947–51.
40. Li S. Testosterone deficiency in aging males: a dual threat to vascular and bone health through coronary artery calcification and osteoporosis—a narrative review. *Aging Male*. 2026;29(1):2662042.
41. Che-Nordin N, Deng M, Griffith JF, Leung JCS, Kwok AWL, Zhu YQ, et al. Prevalent osteoporotic vertebral fractures more likely involve the upper endplate than the lower endplate and even more so in males. *Ann Transl Med*. 2018;6(22):442.
42. Bilek LD, Flores LE, Waltman N, Smith K, Hillstrom D, Yecies L, et al. MON-802 wearable vibration therapy for low bone density: high adherence and usability scores in a 12-month clinical trial. *J Endocr Soc*. 2025;9(Suppl 1):544.
43. Caffarelli C, Tomai Pitinca MD, Al Refaie A, De Vita M, Catapano S, Gonnelli S. Could radiofrequency echographic multispectrometry (REMS) overcome the overestimation in BMD by dual-energy X-ray absorptiometry (DXA) at the lumbar spine? *BMC Musculoskelet Disord*. 2022;23:1–8. Available from: <https://pmc.ncbi.nlm.nih.gov/articles/PMC9118880/>. Accessed on 11 August 2026.
44. Mobasser M, Tarverdizadeh N, Mirghafourvand M, Salehi-Pourmehr H, Ostadrahimi A, Farshbaf-Khalili A. The role of bone turnover markers in screening low bone mineral density and their relationship with fracture risk in the postmenopausal period. *J Res Med Sci*. 2023;28:54.

Cite this article as: Nandivada KKVS, Acharya AS, Jothilingam K, Ahamed SI, John AM, Nandivada V. A DEXA scan at 20 years of age in select subjects may show a z-score "shift to the left" giving time after ethnic corrections to regain optimal peak BMD at 30 years. *Int J Res Orthop* 2026;12:1978-84.